

Patient experience of residents with restricted primary care access during the COVID-19 pandemic

Takuya Aoki ^{1,2}, Yasuki Fujinuma,³ Masato Matsushima¹

To cite: Aoki T, Fujinuma Y, Matsushima M. Patient experience of residents with restricted primary care access during the COVID-19 pandemic. *Fam Med Com Health* 2022;**10**:e001667. doi:10.1136/fmch-2022-001667

ABSTRACT

Objectives To evaluate primary care access for COVID-19 consultation among residents who have a usual source of care (USC) and to examine their associations with patient experience during the pandemic in Japan.

Design Nationwide cross-sectional study.

Setting Japanese general adult population.

Participants 1004 adult residents who have a USC.

Main outcome measures Patient experience assessed by the Japanese version of Primary Care Assessment Tool Short Form (JPCAT-SF).

Results A total of 198 (19.7%) reported restricted primary care access for COVID-19 consultation despite having a USC. After adjustment for possible confounders, restricted primary care access for COVID-19 consultation was negatively associated with the JPCAT-SF total score (adjusted mean difference = -8.61, 95% CI -11.11 to -6.10). In addition, restricted primary care access was significantly associated with a decrease in all JPCAT-SF domain scores.

Conclusions Approximately one-fifth of adult residents who had a USC reported restricted primary care access for COVID-19 consultation during the pandemic in Japan. Our study also found that restricted primary care access for COVID-19 consultation was negatively associated with a wide range of patient experience including first contact. Material, financial and educational support to primary care facilities, the spread of telemedicine and the application of a patient registration system might be necessary to improve access to primary care during a pandemic.

INTRODUCTION

In a context where the COVID-19 pandemic has placed a tremendous burden on health-care systems around the world, primary care capabilities play pivotal roles. Primary care providers aid in the triage and treatment process, help educate patients and render preventive care including vaccination. To bypass physical contact and allow vulnerable patients to have access to primary care from the safety of their homes, some primary care providers leverage telemedicine.¹ In Western countries and Japan, most patients with COVID-19 are seen first by primary care providers because the majority of infected

WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ During the COVID-19 pandemic, several countries reported that medical facilities have refused to provide care for patients with suspected or confirmed COVID-19 infection.

WHAT THIS STUDY ADDS

⇒ Approximately one-fifth of adult residents who had a usual source of care (USC) reported restricted primary care access for COVID-19 consultation. Restricted primary care access for COVID-19 consultation was negatively associated with a wide range of patient experience.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE AND/OR POLICY

⇒ As a health policy issue in Japan, it is necessary to improve access to care by telemedicine expansion. In addition, the free access system and the lack of patient registration with primary care providers may result in residents who have a USC but cannot receive consultation when COVID-19 is suspected.

individuals experience only mild to moderate symptoms.²

In Japan, as in other countries, the primary care sector has performed the initial assessment, including testing for COVID-19, and triaged patients to determine those in need of hospitalisation.³ The government has notified the public that they should first visit their primary care physician (usual source of care (USC)) if they have symptoms suspected of COVID-19, such as fever.⁴ Japanese primary care services are generally provided in both community clinics and outpatient departments of small-sized and medium-sized hospitals that are predominantly privately owned and managed.⁵ The Ministry of Health, Labour and Welfare has recommended that all individuals should have a USC voluntarily⁶ and former surveys have reported that approximately half of the Japanese adult residents had a USC.^{7 8} However, patient registration with primary care physicians has not been institutionalised.



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Division of Clinical Epidemiology, Research Center for Medical Sciences, The Jikei University School of Medicine, Tokyo, Japan

²Section of Clinical Epidemiology, Department of Community Medicine, Graduate School of Medicine, Kyoto University, Kyoto, Japan

³Centre for Family Medicine Development, Japanese Health and Welfare Co-operative Federation, Tokyo, Japan

Correspondence to

Dr Takuya Aoki; taoki@jikei.ac.jp

During the pandemic, several countries reported that medical facilities have refused to provide care for patients with suspected or confirmed COVID-19 infection.⁹ This means that not only residents who do not have a USC but also those who have a USC may have restrictions in accessing COVID-19 care. In Japan, some medical institutions, including primary care facilities, have been rejecting possible COVID-19 patients; therefore, Japan's national and city governments are upping the pressure on medical institutions to take in COVID-19 patients.^{10,11} The refusal may be due to concerns about cluster outbreaks within medical institutions and lack of staff, space for consultation and equipment to deal with the infections. However, it has not been qualified how much residents' access to COVID-19 consultation is restricted owing to their primary care providers' refusal to see patients with suspected COVID-19. Furthermore, whether restricted primary care access during the pandemic is associated with patient experience has not previously been investigated. Patient experience is the core quality measure of patient-centredness, which is globally deemed to be one of the core aims of a healthcare system.^{12,13}

Therefore, in this study, we aimed to evaluate primary care access for COVID-19 consultation among residents who have a USC and to examine their associations with patient experience during the pandemic in Japan.

METHODS

Design, setting and participants

The data for this study were sourced from the National Usual source of Care Survey (NUCS), which was conducted in May 2021 during the COVID-19 fourth wave in Japan. The NUCS was a nationwide mail survey that collected data on the USC, patient experience of primary care, healthcare utilisation, health conditions, health-related quality of life and sociodemographic characteristics of a representative sample of the Japanese adult population.⁷ One of the primary research objectives of the NUCS was to evaluate primary care access for COVID-19 consultation among residents who have a USC and to examine their associations with patient experience. In the NUCS, a nationally representative panel in Japan, which was administered by the Nippon Research Center, was used to select potential participants. This panel comprised approximately 70 000 residents who were selected using a multistage sampling method and participated in a previous survey of the Nippon Research Center.¹⁴ From the panel, 2000 potential participants aged 20–75 years were selected using stratified sampling by age, sex and residential area. The survey participants received ¥500 gift certificates.

Among adult residents who responded to the NUCS, eligible participants in this study were individuals who had a USC. To identify an individual's USC, the following items were used in the Primary Care Assessment Tool (PCAT)¹⁵ and the Medical Expenditure Panel Survey¹⁶: 'Is there a doctor that you usually go to if you

are sick or need advice on your health?' A participant was considered to have a USC if they were able to identify a physician who practices outside university hospitals. According to a previous national survey conducted by the Japan Medical Association in 2020, the proportion of Japanese adults who have a USC was 55.2%.⁸ The eligible participants in this study, who have a USC, were considered to be highly representative because we used the resident panel selected by a probability sampling method.

Measures

Patient experience of primary care

The outcome measure in this study was patient experience of primary care assessed by the Japanese version of Primary Care Assessment Tool Short Form (JPCAT-SF).¹⁷ The JPCAT-SF is based on the PCAT,¹⁵ which was developed by the Johns Hopkins Primary Care Policy Center. This tool is a Japanese version of the PCAT and not a simple Japanese translation of the PCAT. It consists of fewer items than the original version for better usability. A previous study showed that the JPCAT-SF has good reliability and validity.¹⁷ This 13-item tool comprises six multi-item domains addressing the following primary care attributes: first contact, longitudinality, coordination, comprehensiveness (services available), comprehensiveness (services provided) and community orientation.¹⁵ The JPCAT-SF scoring system is structured as follows: each response on a 5-point Likert scale (1=strongly disagree, 2=somewhat disagree, 3=not sure, 4=somewhat agree and 5=strongly agree) is converted into an item score between 0 and 4. The calculated means of item scores in the same domain were multiplied by 25 to yield domain scores ranging from 0 to 100 points. The JPCAT-SF total score is the mean of the six domain scores and reflects an overall measure of primary care experience, with higher scores indicating better patient experience.

Primary care access for COVID-19 consultation

In this study, primary care access for COVID-19 consultation was assessed from the patient's perspective. Restricted primary care access for COVID-19 consultation was defined as impaired access to an individual's USC when COVID-19 was suspected owing to fever during the outbreak. Survey respondents were asked about service availability in their USC using a standardised question 'Please respond to the situation after the outbreak of the new coronavirus. When you have a fever, would someone from your primary care physician's facility see you?' Participants were asked to rate this question on a 4-point scale (1=definitely, 2=probably, 3=probably not and 4=definitely not). We developed this question by modifying the items in the first contact domain of the PCAT and reviewing its face validity. Participants were identified as having restricted primary care access if they responded 'definitely not' or 'probably not' to this question.

Confounding factors

The potential confounding factors were selected on the basis of previous studies that suggested confounding relationships between primary care access and patient experience.^{5 18–20} We assessed the following factors by using a self-administered questionnaire: age, sex, years of education, annual household income, number of chronic conditions, primary care physician location (clinic, hospital, other) and health-related quality of life assessed by the five-level version of the EuroQol five-dimensional questionnaire.²¹ We used a validated list of 20 chronic conditions that were created based on previous multi-morbidity literature and their relevance to the primary care population²²: hypertension, depression/anxiety, chronic musculoskeletal conditions causing pain or limitation, arthritis/rheumatoid arthritis, osteoporosis, chronic respiratory disease (asthma, chronic obstructive pulmonary disease or chronic bronchitis), cardiovascular disease, heart failure, stroke/transient ischaemic attack, stomach problem, colon problem, chronic hepatitis, diabetes, thyroid disorder any cancer in the previous 5 years, kidney disease/failure, chronic urinary problem, dementia/Alzheimer’s disease, hyperlipidaemia and obesity.

Statistical analysis

Descriptive statistics were obtained for the participants’ characteristics and the JPCAT-SF scores. To examine the association between primary care access for COVID-19 consultation and patient experience assessed by the JPCAT-SF total score, we used multivariable linear regression analyses adjusting for confounding factors. In addition, we also performed exploratory analyses to investigate the associations between primary care access and each domain score of the JPCAT-SF using the same models. According to previous studies, a difference of >3-point in patient experience measures linearly scaled to a 0–100 range was considered significant in magnitude with regard to practical importance.^{5 23 24}

For each analysis, we used a two-sided significance level of $p=0.05$. Missing data for independent and dependent variables were adjusted by applying multiple imputations, with 20 imputations, using a fully conditional specification. Statistical analyses were conducted using R, V.4.1.0 (R Foundation for Statistical Computing, Vienna, Austria; www.R-project.org).

RESULTS

Participants’ characteristics

A total of 1757 individuals out of 2000 adult residents responded to the NUCS (response rate: 87.9%). Among them, we excluded 746 respondents who did not have a USC and seven respondents who did not respond to the survey item regarding access to COVID-19 consultation. We performed analyses of the remaining 1004 eligible participants. **Table 1** shows the characteristics of the eligible participants, with or without restricted primary

Table 1 Participants’ characteristics

Characteristic	Total (N=1004)	Primary care access for COVID-19 consultation	
		Not restricted (n=806)	Restricted (n=198)
Age, mean (SD), years	53.1 (15.1)	53.1 (15.4)	53.0 (14.1)
Gender, n (%)			
Female	541 (53.9)	429 (53.2)	112 (56.6)
Education, n (%)			
Less than high school	36 (3.6)	29 (3.6)	7 (3.5)
High school	346 (34.5)	277 (34.4)	69 (34.8)
Junior college	278 (27.7)	224 (27.8)	54 (27.3)
More than or equal to college	320 (31.9)	257 (31.9)	63 (31.8)
Data missing	24 (2.4)	19 (2.4)	5 (2.5)
Annual household income, n (%), million JPY			
<3.00 (=US\$27 000)	168 (16.7)	131 (16.3)	37 (18.7)
3.00–4.99	330 (32.9)	269 (33.4)	61 (30.8)
5.00–6.99	255 (25.4)	201 (24.9)	54 (27.3)
7.00–9.99	165 (16.4)	142 (17.6)	23 (11.6)
≥ 10.00	76 (7.6)	55 (6.8)	21 (10.6)
Data missing	10 (1.0)	8 (1.0)	2 (1.0)
No of chronic conditions, n (%)			
0	322 (32.1)	266 (33.0)	56 (28.3)
1	297 (29.6)	243 (30.1)	54 (27.3)
≥ 2	344 (34.3)	270 (33.5)	74 (37.4)
Data missing	41 (4.1)	27 (3.3)	14 (7.1)
EQ-5D-5L, mean (SD)	0.88 (0.09)	0.89 (0.08)	0.87 (0.12)
Data missing, n (%)	2 (0.2)	2 (0.2)	0 (0.0)
PCP location, n (%)			
Clinic	765 (76.2)	620 (76.9)	145 (73.2)
Hospital	224 (22.3)	173 (21.5)	51 (25.8)
Other	12 (1.2)	11 (1.4)	1 (0.5)
Data missing	3 (0.3)	2 (0.2)	1 (0.5)

EQ-5D-5L, five-level version of the EuroQol five-dimensional questionnaire; PCP, primary care physician.

care access for COVID-19 consultation. Among the eligible participants, 198 (19.7%) reported restricted primary care access for COVID-19 consultation despite having a USC. Compared with participants without restricted primary care access, those with restricted primary care access had more chronic conditions (≥2 chronic conditions, 37.4% vs 33.5%).

Primary care access for COVID-19 consultation and patient experience

Table 2 shows the distribution of JPCAT-SF scores. The average JPCAT-SF total score was 45.9 out of 100 points; the most highly scored domain was longitudinality (54.4), and the most poorly scored domain was first

Table 2 Distribution of JPCAT-SF scores: mean (SD)

	Total		Primary care access for COVID-19 consultation	
	(N=1004)		Not restricted	Restricted
			(n=806)	(n=198)
JPCAT-SF total score	45.9 (17.0)	47.6 (16.8)	38.9 (15.8)	<0.001
JPCAT-SF domain scores				
First contact	32.9 (26.3)	35.4 (27.0)	22.9 (19.9)	<0.001
Longitudinality	54.4 (25.9)	56.1 (25.7)	47.2 (25.9)	<0.001
Coordination	54.5 (29.7)	55.8 (30.0)	49.1 (27.5)	0.004
Comprehensiveness (services available)	49.7 (26.5)	51.8 (26.0)	41.5 (27.2)	<0.001
Comprehensiveness (services provided)	34.4 (31.3)	35.4 (31.7)	30.1 (29.2)	0.034
Community orientation	49.8 (19.6)	51.5 (19.2)	42.9 (19.6)	<0.001

*All scores range from 0 to 100.

†P value by Student's t-test.

JPCAT-SF, Japanese version of Primary Care Assessment Tool Short Form.

contact (32.9). [Table 2](#) also compares the distribution of JPCAT-SF scores with or without restricted primary care access for COVID-19 consultation. The participants with restricted primary care access had lower total scores and all domain scores compared with participants without restricted access.

[Table 3](#) shows the results of the linear regression analyses, examining the association between primary care access to COVID-19 consultation and JPCAT-SF scores as measures of patient experience. After adjustment for possible confounders, restricted primary care access for COVID-19 consultation was negatively associated with the JPCAT-SF total score (adjusted mean difference = -8.61, 95% CI -11.11 to -6.10). In addition, restricted primary care access was significantly associated with a decrease in all JPCAT-SF domain scores. First contact had the strongest association with restricted primary care access

for COVID-19 consultation (adjusted mean difference = -12.26, 95% CI -16.02 to -8.50).

DISCUSSION

This nationwide study in Japan revealed that approximately one-fifth of adult residents who had a USC reported restricted primary care access for COVID-19 consultation owing to their primary care providers' refusal during the pandemic. Our study also found that restricted primary care access for COVID-19 consultation was negatively associated with a wide range of patient experience assessed by the JPCAT-SF. Refusal of COVID-19 care by primary care providers was negatively associated with not only first contact, which is an attribute related to accessibility, but also with other attributes

Table 3 Association between primary care access for COVID-19 consultation and patient experience (n=1004)

Outcome*	Unadjusted		Adjusted†	
	Mean difference (95% CI)	P value	Mean difference (95% CI)	P value
JPCAT-SF total score	-8.68 (-11.26 to -6.11)	<0.001	-8.61 (-11.11 to -6.10)	<0.001
JPCAT-SF domain scores				
First contact	-12.33 (-16.35 to -8.31)	<0.001	-12.26 (-16.02 to -8.50)	<0.001
Longitudinality	-8.90 (-12.89 to -4.91)	<0.001	-8.84 (-12.70 to -4.97)	<0.001
Coordination	-6.73 (-11.37 to -2.10)	0.004	-6.63 (-11.25 to -2.02)	0.005
Comprehensiveness (services available)	-10.40 (-14.51 to -6.28)	<0.001	-10.34 (-14.48 to -6.20)	<0.001
Comprehensiveness (services provided)	-5.27 (-10.13 to -0.42)	0.034	-5.21 (-10.09 to -0.33)	0.037
Community orientation	-8.48 (-11.48 to -5.48)	<0.001	-8.34 (-11.38 to -5.31)	<0.001

Reference group: Without restricted primary care access.

*All scores range from 0 to 100.

†Adjusted for age, sex, years of education, annual household income, number of chronic conditions, EQ-5D-5L and primary care physician location.

EQ-5D-5L, five-level version of EuroQol five-dimensional questionnaire; JPCAT-SF, Japanese version of Primary Care Assessment Tool Short Form.

including longitudinality, coordination, comprehensiveness and community orientation.

Although there have been domestic and international news reports on the refusal of patients with suspected or confirmed COVID-19 in primary care facilities, few studies have presented specific data. In a previous simulated patient study in primary care facilities in the USA, 20% of participating facilities guided patients with suspected COVID-19 to consultation in an emergency department or COVID-19 hotline.²⁵ Such a provider's response to COVID-19 may have a broad impact on the residents' experience because COVID-19 is a health issue of great concern to residents. Since primary care attributes such as first contact, longitudinality, coordination, comprehensiveness and community orientation play an important role in COVID-19 care,²⁶ the refusal of patients with suspected COVID-19 may have led to residents' evaluation of their primary care provider as lacking in these characteristics. This overall poor primary care experience may lead to poor adherence to treatment of both communicable and noncommunicable diseases, preventive care and inefficient patient behaviours such as bypassing a USC to seek care at higher-level healthcare facilities.^{27–30}

Our study indicated that timely access to primary care during a pandemic may be a critical issue in improving patient experience. Refusal of COVID-19 patients is not only an issue at the level of medical providers and facilities, but also at the healthcare system. For primary care facilities to respond to the pandemic, they need support from the government and outside medical institutions, such as education on infectious disease care and control, supplying necessary supplies and financial support. As a health policy issue in Japan, the application and spread of telemedicine have been slow even during the pandemic.³¹ Therefore, it is necessary to improve access to care by telemedicine expansion while considering the quality, safety and equity of care, especially for vulnerable people who have multimorbidity. In addition, the free access system and the lack of patient registration with primary care providers may lead some primary care providers not to responsibly manage the health of individual residents, including when COVID-19 is suspected.

To the best of our knowledge, this is the first study to report residents' primary care access for COVID-19 consultation and examine their associations with patient experience in Japan. A key strength of our study is the use of data from a nationwide study, with a sample representative of the Japanese adult population, which allows for generalisation of its results to the wider population. Another strength is the high study response rate compared with other national surveys. The PCAT is a validated and internationally established tool for evaluating patient experience of primary care attributes.

Our study also has several potential limitations. First, our assessment of primary care access was based on residents' perspectives and not on objective data from providers. However, subjective assessment of accessibility is also crucial because it directly relates to healthcare utilisation of residents.³² Second, we collected information on access to

consultation when limited to fever, a typical symptom of COVID-19, but not for other symptoms and conditions already confirmed as COVID-19. Third, given that the data were cross-sectional, a causal relationship between restricted primary care access and patient experience can definitively not be established. Fourth, this study was conducted in Japan, which does not have a patient registration system for primary care providers, and this should be considered when generalising the results of this study to other countries.

CONCLUSIONS

Approximately one-fifth of adult residents who had a USC reported restricted primary care access for COVID-19 consultation during the pandemic in Japan. Our study also found that restricted primary care access for COVID-19 consultation was negatively associated with a wide range of patient experience including first contact. Material, financial and educational support to primary care facilities, the spread of telemedicine, and the application of a patient registration system might be necessary to improve access to primary care during a pandemic.

Contributors All authors (TA, YF and MM) of the paper contributed the conception or design of the work. TA performed the statistical analyses. TA, YF and MM interpreted the analyses. TA drafted the manuscript. All authors reviewed and edited the manuscript, contributed to the discussion of the data and performed critical review of the manuscript. All authors gave the final approval of the manuscript before submission. TA is the guarantor of the work and accepts full responsibility for the presented content.

Funding This work was supported by JSPS KAKENHI Grant Number JP20K18849.

Competing interests TA reports grants from JSPS KAKENHI, during the conduct of the study; and TA received lecture fees and lecture travel fees from the Centre for Family Medicine Development of Japanese Health and Welfare Co-operative Federation. MM received lecture fees and lecture travel fees from the Centre for Family Medicine Development of Japanese Health and Welfare Co-operative Federation. MM's son-in-law worked at IQVIA Services Japan K.K. which is a contract research organisation and a contract sales organisation. MM's son-in-law works at SYNEOS HEALTH CLINICAL K.K. which is a contract research organisation and a contract sales organisation.

Patient consent for publication Consent obtained directly from patient(s).

Ethics approval The institutional review board of the Jikei University School of Medicine approved this study (approval no. 32-416(10505)). Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement No data are available. Due to the nature of this research, participants of this study did not agree for their data to be shared publicly, so supporting data are not available.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Takuya Aoki <http://orcid.org/0000-0002-8232-2155>

REFERENCES

- Lee JQ, Loke W, Ng QX. The role of family physicians in a pandemic: a blueprint. *Health Care* 2020;8:198.



- 2 Kearon J, Risdon C. The role of primary care in a pandemic: reflections during the COVID-19 pandemic in Canada. *J Prim Care Community Health* 2020;11:215013272096287.
- 3 Haruta J, Horiguchi S, Miyachi J, *et al.* Primary care physicians' narratives on COVID-19 responses in Japan: professional roles evoked under a pandemic. *J Gen Fam Med* 2021. doi:10.1002/jgf2.452. [Epub ahead of print: 14 May 2021].
- 4 Ministry of Health, Labour and Welfare. Medical institutions for treatment and testing (Japanese). Available: https://www.mhlw.go.jp/stf/seisakunitsuite/bunya/kenkou_iryuu/covid19-jyushinsoudancenter.html [Accessed 7 Oct 2021].
- 5 Aoki T, Yamamoto Y, Fukuhara S. Comparison of primary care experience in hospital-based practices and community-based office practices in Japan. *Ann Fam Med* 2020;18:24–9.
- 6 Ministry of Health, Labour and Welfare. Report on clarification and coordination of outpatient functions, strengthening of kakaritukei functions (Japanese). Available: <https://www.mhlw.go.jp/content/10801000/000704605.pdf> [Accessed 12 May 2022].
- 7 Aoki T, Fujinuma Y, Matsushima M. Usual source of primary care and preventive care measures in the COVID-19 pandemic: a nationwide cross-sectional study in Japan. *BMJ Open* 2022;12:e057418.
- 8 The Japan Medical Association Research Institute. The 7th survey on attitudes toward medical care in Japan (Japanese). Available: <https://www.jmari.med.or.jp/download/WP448.pdf> [Accessed 12 May 2022].
- 9 David Williams O, Yung KC, Grépin KA. The failure of private health services: COVID-19 induced crises in low- and middle-income country (LMIC) health systems. *Glob Public Health* 2021;16:1320–33.
- 10 The Japan Times. Hospitals in Japan refusing to test many who suspect they have COVID-19. Available: <https://www.japantimes.co.jp/news/2020/02/26/national/hospitals-refuse-coronavirus-patients/> [Accessed 12 May 2022].
- 11 This Week in Asia. Available: https://www.scmp.com/week-asia/health-environment/article/3146172/japan-name-and-shame-hospitals-refuse-covid-19?module=perpetual_scroll&pgtype=article&campaign=3146172 [Accessed 12 May 2022].
- 12 Institute of Medicine. *Committee on quality of health care in America. crossing the quality chasm: a new health system for the 21st century.* Washington, DC: National Academies Press, 2001.
- 13 Berwick DM, Nolan TW, Whittington J. The triple aim: care, health, and cost. *Health Aff* 2008;27:759–69.
- 14 Nippon Research Center. Nippon research center. Available: <https://www.nrc.co.jp/english/index.html> [Accessed 12 May 2022].
- 15 Shi L, Starfield B, Xu J. Validating the adult primary care assessment tool. *J Fam Pract* 2001;50:161–75.
- 16 Agency for Healthcare Research and Quality. Medical expenditure panel survey. Available: <https://www.meps.ahrq.gov/mepsweb/index.jsp> [Accessed 12 May 2022].
- 17 Aoki T, Fukuhara S, Yamamoto Y. Development and validation of a Concise scale for assessing patient experience of primary care for adults in Japan. *Fam Pract* 2020;37:137–42.
- 18 Levesque J-F, Harris MF, Russell G. Patient-centred access to health care: conceptualising access at the interface of health systems and populations. *Int J Equity Health* 2013;12:18.
- 19 Campbell JL, Ramsay J, Green J. Age, gender, socioeconomic, and ethnic differences in patients' assessments of primary health care. *Qual Health Care* 2001;10:90–5.
- 20 Lyrtzopoulos G, Elliott M, Barbieri JM, *et al.* Understanding ethnic and other socio-demographic differences in patient experience of primary care: evidence from the English general practice patient survey. *BMJ Qual Saf* 2012;21:21–9.
- 21 Herdman M, Gudex C, Lloyd A, *et al.* Development and preliminary testing of the new five-level version of EQ-5D (EQ-5D-5L). *Qual Life Res* 2011;20:1727–36.
- 22 Fortin M, Almirall J, Nicholson K. Development of a research tool to document self-reported chronic conditions in primary care. *J Comorb* 2017;7:117–23.
- 23 Paddison CAM, Elliott MN, Haviland AM, *et al.* Experiences of care among medicare beneficiaries with ESRD: Medicare consumer assessment of healthcare providers and systems (CAHPS) survey results. *Am J Kidney Dis* 2013;61:440–9.
- 24 Warren FC, Abel G, Lyrtzopoulos G, *et al.* Characteristics of service users and provider organisations associated with experience of out of hours general practitioner care in England: population based cross sectional postal questionnaire survey. *BMJ* 2015;350:1–9.
- 25 Kyle MA, Tipirneni R, Thakore N, *et al.* Primary care access during the COVID-19 pandemic: a simulated patient study. *J Gen Intern Med* 2021;36:3766–71.
- 26 Henry TL, Rich EC, Bazemore A. Comprehensiveness—the need to Resurrect a sagging Pillar of primary care. *J Gen Intern Med* 2022;37:229–31.
- 27 Doyle C, Lennox L, Bell D. A systematic review of evidence on the links between patient experience and clinical safety and effectiveness. *BMJ Open* 2013;3:e001570.
- 28 Aoki T, Inoue M. Primary care patient experience and cancer screening uptake among women: an exploratory cross-sectional study in a Japanese population. *Asia Pac Fam Med* 2017;16:3.
- 29 Kaneko M, Aoki T, Goto R, *et al.* Better patient experience is associated with better vaccine uptake in older adults: multicentered cross-sectional study. *J Gen Intern Med* 2020;35:3485–91.
- 30 Aoki T, Yamamoto Y, Ikenoue T, *et al.* Effect of patient experience on bypassing a primary care gatekeeper: a multicenter prospective cohort study in Japan. *J Gen Intern Med* 2018;33:722–8.
- 31 Miyawaki A, Tabuchi T, Ong MK, *et al.* Age and social disparities in the use of telemedicine during the COVID-19 pandemic in Japan: cross-sectional study. *J Med Internet Res* 2021;23:e27982.
- 32 Haggerty JL, Lévesque J-F, Santor DA, *et al.* Accessibility from the patient perspective: comparison of primary healthcare evaluation instruments. *Healthc Policy* 2011;7:94–107.