

Female-to-male sexual transmission of Zika virus infection

Beuy Joob,¹ Viroj Wiwanitkit²

To cite: Joob B, Wiwanitkit V. Female-to-male sexual transmission of Zika virus infection. *Fam Med Com Health* 2019;7:e000183. doi:10.1136/fmch-2019-000183

Received 25 May 2019
Revised 12 June 2019
Accepted 17 June 2019



© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Sanitation 1, Medical Academic Center, Bangkok, Thailand

²Department of Community Medicine, Dr DY Patil Vidyapeeth University, Pune, India

Correspondence to

Dr Beuy Joob;
beuyjoob@hotmail.com

Dear Editor,

We read the publication on 'The challenge given by Zika virus' with a great interest.¹ Rodriguez and Sebastian mentioned that 'No sexual transmission of Zika virus from infected women to their partners and from infected people without symptoms has been reported'.¹ Indeed, sexual transmission is a possible mode of Zika virus transmission.² Either homosexual or heterosexual contact might be the starting point of pathogenic virus transmission.³ Regarding female-to-male sexual transmission of Zika virus, there are some reports on this problem. Davidson *et al* first reported a suspected female-to-male sexual transmission of Zika Virus from New York city in 2016.⁴ This case report had several laboratory evidences showing the possibility of disease transmission.⁴ After that, there are many reports on this issue.⁵ As noted by Sherley and Ong, 'Unusually for a mosquito-borne virus, sexual spread has also been reported; with cases of male-to-female, female-to-male and male-to-male sexual transmission all now published in the scientific literature'.⁵ According to the summative analysis by Moreira *et al* on sexual transmission of Zika virus, female-to-male transmission is detected in 3.7%.⁶ Nevertheless, the common difficulty for diagnosis of a male-to-female or female-to-male sexual transmission of Zika virus is the ruling out of the chance of mosquito transmission, mosquito bite, of the two partners.⁷ The diagnosis requires both

clinical epidemiological molecular laboratory investigations for final confirmation.⁷

Contributors Both authors have contributed equally.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; internally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

1. Vargas Rodriguez JS, Rodriguez V, Sebastian J. The challenge given by Zika virus. *Fam Med Community Health* 2018;6:208–10.
2. Joob B, Wiwanitkit V. Sexual transmission of Zika virus: what is the issue for awareness? *Int Marit Health* 2016;67.
3. Sookaromdee P, Wiwanitkit V. Homosexual and Zika virus infection: is it a new start of new problem for gay? *Indian J Sex Transm Dis AIDS* 2018;39:61–2.
4. Davidson A, Slavinski S, Komoto K, *et al*. Suspected female-to-male sexual transmission of Zika virus. *MMWR Morb Mortal Wkly Rep* 2016;65:716–7.
5. Sherley M, Ong C-W. Sexual transmission of Zika Virus: a literature review. *Sex Health* 2018;15:183–99.
6. Moreira J, Peixoto TM, Siqueira AM, *et al*. Sexually acquired Zika Virus: a systematic review. *Clin Microbiol Infect* 2017;23:296–305.
7. Wiwanitkit S, Wiwanitkit V. Sexually transmitted Zika virus infection: a new tropical disease. *Asian Pac J Trop Dis* 2016;6:757–8.