

# Care and social support from family and community in patients with pulmonary tuberculosis in Pakistan

Shahab E Saqib ,<sup>1</sup> Mokbul Morshed Ahmad,<sup>1</sup> Sanaullah Panezai<sup>2</sup>

**To cite:** Saqib SE, Ahmad MM, Panezai S. Care and social support from family and community in patients with pulmonary tuberculosis in Pakistan. *Fam Med Com Health* 2019;7:e000121. doi:10.1136/fmch-2019-000121  
Accepted

## ABSTRACT

**Objectives** This study aimed to validate the Medical Outcome Study–Social Support Survey (MOS-SSS) instrument in Pakistan and to quantify the nature of care and social support currently extended to patients with pulmonary tuberculosis (TB) in Pakistan.

**Design** This is a cross-sectional study design conducted within a period of 3 months from 1 November 2016 to 31 January 2017.

**Participants** A total of 269 patients, registered at 11 TB centres and private clinics, were interviewed through an interviewer-administered questionnaire.

**Main outcome measures** Cronbach's alpha was used to measure the internal consistency and reliability of the MOS-SSS survey instrument. Univariate and multivariable logistic regressions have been used to explore the association between care and social support, and socioeconomic factors.

**Result** This study validated the use of the MOS-SSS among patients with TB in Pakistan. Findings of the study revealed significant differences among the socioeconomic groups of patients in each subscale of social support. Additionally, results of logistic regressions showed that patients who were older (adjusted OR=6.17, 95% CI 1.55 to 24.59,  $p \leq 0.01$ ), male (adjusted OR=2.73, 95% CI 1.49 to 4.98,  $p \leq 0.01$ ), widow (adjusted OR=0.17, 95% CI 0.04 to 0.80,  $p \leq 0.05$ ), and had a larger household size (adjusted OR=5.69, 95% CI 1.32 to 24.65,  $p \leq 0.05$ ), higher monthly income (adjusted OR=2.00, 95% CI 1.11 to 3.60,  $p \leq 0.05$ ) and house ownership (adjusted OR=1.99, 95% CI 1.10 to 3.60,  $p \leq 0.05$ ) were significant factors associated with the extent of care and social support that the participants received.

**Conclusion** To cure TB, this study suggests a coordinated approach that includes not only clinical services to address this issue but also a strong social support system based on family and community necessary throughout the treatment process.

## INTRODUCTION

Tuberculosis (TB) is considered a major issue of public health with an estimated 10.4 million new TB cases worldwide.<sup>1</sup> Among these cases, 5.9 million cases are reported to be among men, 3.5 million among women, 1.0 million among children and 2.1 million in individuals living with HIV, accounting for all new TB cases.<sup>1</sup>

## Key points

- ▶ This study validates the use of the Medical Outcome Study–Social Support Survey among patients with tuberculosis (TB) in Pakistan.
- ▶ Results revealed that social support received by patients with TB from families and communities was limited.
- ▶ Findings suggest that patients with TB need social care and support from families and communities, along with standard medical treatment.

Besides, along with HIV/AIDS, TB is a major cause of mortality worldwide and places the heaviest burden on the poorest and most vulnerable population of the society.<sup>2</sup> In South Asia, Pakistan has the highest incidence rate (276/100 000) between 2006 and 2014.<sup>3</sup> Pakistan ranks fifth among the high burden countries and accounts for about 61% of the TB burden in the East Mediterranean region. Each year, approximately 0.42 million new cases of TB emerge, among which half of the cases are sputum smear positive. Pakistan is also estimated to have the fourth highest prevalence of multidrug-resistant TB globally.<sup>4</sup>

This slow rate of decline in TB shows that a biomedical approach alone will not be sufficient for ending the TB epidemic and achieving its elimination targets. There is an increasing consensus that social determinants of TB play a vital role in fighting against this disease.<sup>5</sup> Despite well-established treatment regimens, TB remains a burden to public health; it affects both poor and marginalised populations who may not have access to social support, including migrants and the homeless.<sup>6</sup> If patients are carefully treated and observed by family and the surrounding community, this may result in increased rates of compliance to the treatment and enhanced psychological well-being of the patient.<sup>7</sup> Conversely, in cases where compliance to TB therapy does not



© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

<sup>1</sup>Department of Regional and Rural Development Planning, Asian Institute of Technology, Bangkok, Thailand

<sup>2</sup>Department of Geography, University of Balochistan, Quetta, Pakistan

## Correspondence to

Dr Shahab E Saqib;  
shahabmomand@gmail.com

occur as with many cases today, this can lead to drug resistance, prolonged infection and death.<sup>8</sup> For the empirical management of TB, a study revealed that the major drivers of drug-resistant TB are inappropriate drug regimens, insufficient attention to TB, and inadequate care and social support for patients in facilitating full completion of the administered treatment.<sup>9</sup>

As a condition, TB has been linked to several negative social outcomes for patients. For instance, due to the contagious nature of TB, people avoid being in close proximity to individuals infected with TB and think of themselves to be infected. Additionally, during the initial stages of active TB, those under medical care are required to remain physically isolated for 2 weeks or longer, receiving treatment to ensure that they are no longer contagious.<sup>10</sup> However, whether individuals are contagious or not, the stigma originally caused by this ill effect still remains. Within the community, individuals do not forget when someone close carries a life-threatening disease. Patients with TB are often said to be silently isolated by the community, and contact is observed to be avoided with patients with TB. For instance, people are refusing to drink tea offered by patients in social gatherings and not greeting patients physically when they meet.<sup>11</sup> The factors of isolation and other social consequences are described as significant in delaying the diagnosis in TB, especially among patients who are female.<sup>11</sup> For instance, in Thailand, 65% of patients are found to be highly stigmatised by TB.<sup>12</sup> Therefore, due to these different consequences, TB has multidimensional impacts in both the individual and domestic spheres.

Research shows that social integration and social support are beneficial for good health.<sup>13</sup> Patients with TB have several medical and non-medical needs that should be fulfilled to cure this disease. If the patients are not carefully treated and observed by family and community, it may lead to non-compliance of treatment. Non-compliance to TB therapy can lead to drug resistance, prolonged infectiousness and death.<sup>8</sup> A meta-analysis study found that social support and social relationship influence risk of mortality.<sup>14</sup> Furthermore, it is also reported that stronger social relationships increase the likelihood of survival by 50% as compared with those who had no social relationships. Odds ratios were 1.9 for social integration, 1.5 for social networks, 1.4 for perceived social support and 1.2 for received social support.<sup>14</sup> In Pakistan, a study is conducted on the socioeconomic consequences of TB.<sup>15</sup> Another study has been carried out on the need for a holistic approach to address all aspects of TB.<sup>16</sup> While acknowledging the importance of combining non-medical care with medical treatment, previous studies have assessed social support either by quantifying social relationships<sup>17–19</sup> or by other qualitative aspects.<sup>20 21</sup> Limited studies have explicitly assessed the care and social support provided to patients with TB within the context of Pakistan, and no previous studies, in spite of their contributions, have

explicitly assessed the social support received by patients with TB. Therefore, this study attempts primarily to validate the Medical Outcome Study–Social Support Survey (MOS-SSS) instrument in Pakistan. Second, this is our first attempt to quantify the nature of care and social support currently extended to patients with TB in Khyber Pakhtunkhwa, Pakistan.

## METHODOLOGY

### Study design

This is a cross-sectional quantitative study that is designed to test the reliability of MOS-SSS in Pakistan and explores the relationship with other socioeconomic and demographic factors.

### Study duration

The study was conducted within a period of 3 months from 1 November 2016 to 31 January 2017.

### Study area

The study was conducted in District Mardan, Khyber Pakhtunkhwa Province of Pakistan. Mardan is the second largest city of the province and 19th in Pakistan.<sup>22</sup> This district is administratively divided into three subdistricts: Mardan, Takht Bhai and Katlang. The total population of the district is 1.46 million, of which 0.75 million are male and 0.71 million are female.<sup>23</sup> We purposively selected Khyber Pakhtunkhwa due to its high prevalence of TB cases, where approximately 58 449 of new cases of TB were reported in 2014.<sup>24</sup> In response to the high prevalence, the provincial government has considered this a serious issue and passed an act in 2016 known as ‘the KP TB Notification Bill, 2016’. This act stipulated TB as a disease to be notified by all involved stakeholders, including medical practitioners, private and government clinics and community leaders.<sup>25</sup>

### Study population and sampling

A multistage sampling was adopted to select the location of the study and the participants. A total of 5624 new patients with TB were registered at 11 TB centres and private clinics in the district in 2016.<sup>26</sup> We excluded the first quarter and second quarter registered patients, extrapulmonary cases, children and the significantly elderly within the target population due to their inability to participate in the interviews. Interviews were conducted in clinics. We have used the formula of a previous study and a sample size of 280 was calculated.<sup>27</sup> The total sample size was proportionally allocated among male and female participants, where male patients were 132 and female patients were 148. Lastly, those patients who were under treatment for at least 1 month in the last two quarters of year 2016 were randomly selected. However, in the third month of data collection, due to sociocultural constraints and incomplete questionnaire, the total sample size declined to 269 and consisted of 130 male and 139 female respondents.

### Inclusion and exclusion criteria

This study employed both TB smear-positive and smear-negative patients under treatment for at least the previous 4 weeks, and at the time of the interview, they had at least two visits to the TB centre. We excluded the extrapulmonary patients from the study population. The patients below age 15 and above 60 were also excluded from the sample population because the children and the very old age patients could not answer the questions.

### Study instrument

The MOS-SSS is a 19-item scale survey that contains subscales that include the domains of emotional/informational, tangible, affectionate, positive social interaction, additional support and one additional item.<sup>19 28</sup> The MOS-SSS is one of the most widely used instruments and is validated in different cultures, languages and contexts.<sup>29–32</sup> Data were collected on a Likert scale and were transformed from 0 to 100 for each subscale, where the higher values imply higher levels of care and social support. We obtained a score for each subscale and calculated the mean scores for each item in the subscale. The questionnaire was translated into Pushto language by following the WHO guidelines.<sup>33</sup> Cronbach's alpha was used to measure the internal consistency and reliability. The questionnaire also contained demographic and socioeconomic information of the patients, such as age, sex, marital status, location of the house, income and main occupation.

### Statistical analysis

Statistical tests such as t-test and analysis of variance were used to see the differences in subscale items along with socioeconomic groups. This study is further extended to explore the relationship between social support and other socioeconomic characteristics of the patients. Therefore, we used univariate logistic and multivariable logistic regressions. Furthermore, the social support index was converted into a dichotomous variable, taking the mean value as a cut point following other relevant studies.<sup>34–37</sup>

## RESULTS

### Demographic information of the participants

Patients were mostly young (56.13% were of ages ranging from 15 to 30 years). Among the total participants, 51.67% were female. Married patients were 60.2%, and 54.65% had 5–10 household members. Majority of them were illiterate (51.28%). In addition, most of the patients (62.45%) were living in rural areas (table 1). Most of the patients were from the low-income class: 54.65% made <US\$250 per month as household income. Besides, 34.20% and 28.63% were

**Table 1** Demographic information of participants

Demographic information	Frequency	Percentage (%)
<b>Age groups</b>		
15–20	63	23.42
21–30	88	32.71
31–40	54	20.10
41–50	45	16.72
51–60	19	7.06
<b>Sex</b>		
Female	139	51.67
Male	130	48.33
<b>Marital status</b>		
Unmarried	91	33.80
Married	162	60.20
Widowed	16	6.00
<b>Household size</b>		
1–5	72	26.77
5–10	147	54.65
11–15	32	11.90
>15	18	6.69
<b>Location</b>		
Urban	101	37.55
Rural	168	62.45
<b>Literacy</b>		
Illiterate	146	54.28
Literate	123	45.72
<b>Monthly income</b>		
<250	147	54.65
250–500	83	30.86
500–750	23	8.55
>750	16	5.94
<b>House ownership</b>		
No	151	56.13
Yes	118	43.87
<b>Main households' occupation</b>		
Agriculture	77	28.63
Trader	41	15.24
Labour	92	34.20
Others	26	9.66
Formal Job	33	12.27

wage labourers and had agriculture as the main source of the family's income, respectively.

### Summary of social support items

In all the eight items of subscales, most of the respondents had an emotional/information support score of 3.5. The

**Table 2** Frequency distribution of social support measuring items

Support item	None of the time (1)	A little of the time (2)	Some of the time (3)	Most of the time (4)	All of the time (5)	Mean	SD
<b>Emotional/information support</b>							
<b>Availability of someone</b>							
To listen to you	16 (5.95)	36 (13.38)	66 (24.54)	96 (35.69)	55 (20.45)	3.51	1.13
To give information	8 (2.97)	43 (15.99)	71 (26.39)	96 (35.69)	51 (18.96)	3.52	1.06
To give you advice	5 (1.86)	39 (14.50)	73 (27.14)	106 (39.41)	46 (17.10)	3.55	0.99
To talk about yourself or your problems	11 (4.09)	42 (15.61)	68 (25.28)	96 (35.69)	52 (19.33)	3.51	1.09
Whose advice you want	9 (3.35)	45 (16.73)	79 (29.37)	92 (34.20)	44 (16.36)	3.43	1.05
To share your private worries	5 (1.86)	46 (17.10)	71 (26.39)	103 (38.29)	44 (16.36)	3.50	1.01
Turn to for suggestions	13 (4.83)	41 (15.24)	75 (27.88)	88 (32.71)	52 (19.33)	3.46	1.11
To understand your problems	10 (3.72)	43 (15.99)	76 (28.25)	94 (34.94)	46 (17.10)	3.46	1.06
<b>Total</b>						3.5	0.02
<b>Tangible support</b>							
To help if you were confined to bed	9 (3.35)	57 (21.19)	50 (18.59)	98 (36.43)	55 (20.45)	3.49	1.34
Take you to the doctor	8 (2.97)	44 (16.36)	59 (21.93)	114 (42.38)	44 (16.36)	3.53	1.03
To prepare your meals	6 (2.23)	50 (18.59)	49 (18.22)	118 (43.87)	46 (17.10)	3.55	1.04
Help with daily chores	6 (2.23)	50 (18.59)	52 (19.33)	114 (42.38)	47 (17.47)	3.54	1.05
<b>Total</b>						3.53	0.01
<b>Affectionate support</b>							
Shows you love and affection	15 (5.58)	49 (18.22)	67 (24.91)	111 (41.26)	27 (10.04)	3.32	1.06
Loves you	19 (7.06)	42 (15.61)	80 (29.74)	87 (32.34)	41 (15.24)	3.33	1.23
Cares for you	15 (5.58)	49 (18.22)	77 (28.62)	95 (35.32)	33 (12.27)	3.30	1.78
<b>Total</b>						3.31	0.01
<b>Positive social interaction</b>							
Have a good time with	9 (3.35)	50 (18.59)	108 (40.15)	82 (30.48)	20 (7.43)	3.20	0.94
Get together for relaxation	1 (0.37)	82 (30.48)	77 (28.62)	95 (35.32)	14 (5.20)	3.14	0.93
Do something enjoyable with	4 (1.49)	56 (20.82)	106 (39.41)	89 (33.09)	14 (5.20)	3.20	0.88
<b>Total</b>						3.12	0.02
<b>Overall total</b>						3.42	0.03
<b>Additional item</b>							
Help you get your mind off things	13 (4.83)	45 (16.73)	125 (46.47)	62 (23.05)	24 (8.92)	3.14	0.96

Note: Figures in parentheses are percentages.

mean score of all items was 3.42 (table 2). 'To give you advice' and 'to prepare your meals' had the highest score (3.55). In contrast, 'get together for relaxation' and 'help you get your mind off things' showed the lowest score (3.14).

#### Descriptive statistics of subscale measuring items

The mean scores of the subscales are from 53.62 (minimum) to 63.22 (maximum). Cronbach's alpha values ranged from 0.79 to 0.95. The overall average value of social support was 60.14 and was taken as a cut point to make a dichotomous dependent variable (table 3).

#### Comparison among patients' social support scores in different subscales

As shown in table 4, there is a significant difference ( $p \leq 0.01$ ) in almost all subscales of social support with all age groups except for tangible support. The results for overall support index was higher (66.31) in men than in the women (54.37), and there is a significant difference ( $p \leq 0.01$ ) between men and women in all subscales of social support. Unmarried patients received better emotional/information support than other patients who were married and widows ( $p \leq 0.01$ ). Patients with a larger household's size or owned the house receive better social support in all aspects (all  $p$  values were  $\leq 0.05$ ). Patients

**Table 3** Mean scores of the subscales

Measures	K	Mean	DS	Range	Alpha
Emotional/information support	8	62.34	23.05	0–100	0.95
Tangible support	4	63.22	24.58	0–100	0.93
Affectionate support	3	57.96	23.67	0–100	0.84
Positive social interaction	3	54.52	19.21	0–100	0.82
Additional item	1	53.62	24.11	0–100	0.79
Overall support	19	60.14	17.02	0–100	

who were living in urban areas received better tangible support ( $p \leq 0.05$ ), and literate patients had better positive interaction ( $p \leq 0.05$ ). There was no significant difference in all social support scales among the patients' main households' occupation groups.

### Results of the regression model

In regression analysis, the *p* value of the Hosmer and Lemeshow test was 0.487, revealing good model fit (table 5). Factors such as the age of the patients, sex, marital status, household size, monthly income and house ownership were identified as significant factors associated with the social support received by the patients. There was no significant difference in main households' occupation. Therefore, we have excluded this variable in the regression model. Patients in the age group 51–60 years were more likely to receive social support than the adolescent group between 15 and 20 years old (adjusted OR=6.17, 95% CI 1.55 to 24.59). Men were more likely to receive social support than women (adjusted OR=2.73, 95% CI 1.49 to 4.98). Social support received by widows was lower than unmarried participants (adjusted OR=0.17, 95% CI 0.04 to 0.80). Patients who had >15 family members received a higher level of social support than the lowest category of the household size (adjusted OR=5.69, 95% CI 1.32 to 24.65). Patients with >US\$750 monthly income were more likely to acquire social support than the lowest income group (adjusted OR=2.0, 95% CI 1.11 to 3.60). Patients who owned their houses were also more likely to get support than non-owners (adjusted OR=1.99, 95% CI 1.10 to 3.60).

### DISCUSSION

TB targets most often the utmost vulnerable in any population such as the poor, malnourished and homeless who have no access to social support.<sup>6</sup> Therefore, the importance of the care and social support to patients with TB from family and community can never be neglected during treatment. The findings of this study revealed that MOS-SSS is a valid instrument used for measuring the levels of social support provided to patients with TB in Pakistan by substantiating high reliability in all the subscale measuring items.<sup>19</sup> Patients with pulmonary TB in our study received limited social support. However, it was still higher than another study conducted in China (60.14 vs 34.56).<sup>38</sup>

The score of tangible support indicate that respondents received a good level of support from their families, friends and the surrounding community. However, the results on affectionate support and positive social interaction showed that in these subscales, the patients received limited social support.

The findings of the current study revealed that older patients received a higher care and social support score than young patients. These findings are different from the study in China, which demonstrated that young patients received lower social support (27.75) than elderly patients (31.60), and the population between 40 and 59 years old showed the highest score (33.60).<sup>38</sup> Young patients usually have more energy and financial power. Therefore, they reported less social support from families and community. The elderly were perceived to require more attention from family, friends and the community. Therefore, they reported higher social support. Besides, in the cultural and religious contexts of Pakistan, supporting the sick and the elderly is believed to be rewarded later by God: 'Helping the one who is sick and serving him is an act of charity'. The Prophet (blessings and peace of Allah be on him) said, 'Helping a man onto his mount or lifting up his luggage onto, it is a charity' (narrated by a Muslim, 1009). The findings of the study are consistent with a study conducted among Indians in America. They revealed that the respect for elders is an important traditional value that may translate into higher social support in American Indians.<sup>39</sup>

Additionally, the present study revealed that male patients received higher levels of social support than female patients. As Pakistani society is patriarchal, men are perceived as the primary authority figures in the home, and men hold a superior position in relation to women in both the financial and social spheres.<sup>40</sup> Male and female patients with TB in Pakistan face many social and economic problems, but female patients suffered more than men.<sup>15</sup> A study reported that female patients with TB had less social support from their families.<sup>16</sup> This result is in contrast with the findings of a study from China, which revealed no difference between men and women receiving the social support.<sup>38</sup>

In all subscales, marital status is an important determinant of social support. Widows showed a negative association with social support, which implies that widows receive less social support than their unmarried and married

**Table 4** Socioeconomic characteristics and patients' social support scores in all subscales

Socioeconomic characteristics	Emotional/information support	Tangible support	Positive interaction	Affectionate	Overall support index
Age (years)					
15–20	50.65 (25.80)	61.81 (27.77)	48.55 (22.58)	47.88 (19.85)	52.21 (18.14)
21–30	62.79 (21.03)	63.64 (23.57)	58.62 (20.51)	54.07 (17.05)	60.32 (14.21)
31–40	66.84 (20.33)	61.81 (24.75)	58.33 (24.66)	57.72 (17.72)	62.28 (15.87)
41–50	66.53 (21.57)	64.44 (22.84)	64.44 (25.28)	57.78 (20.90)	63.98 (17.99)
51–60	76.32 (18.23)	67.11 (23.28)	69.74 (25.19)	61.84 (21.75)	70.43 (16.93)
P value†	p≤0.01**	0.91	p≤0.01**	p≤0.01†	p≤0.01**
Sex					
Female	55.62 (22.90)	59.44 (24.85)	52.16 (24.72)	48.56 (20.29)	54.37 (17.13)
Male	69.52 (21.02)	67.26 (23.73)	64.17 (20.85)	60.90 (15.73)	66.31 (14.61)
P value	p≤0.01**	p≤0.01**	p≤0.01**	p≤0.01**	p≤0.01**
Marital status					
Unmarried	63.94 (20.29)	61.13 (23.60)	55.86 (24.95)	54.94 (18.21)	60.06 (16.30)
Married	63.06 (23.69)	64.74 (24.10)	59.72 (22.85)	54.63 (19.59)	61.10 (17.09)
Widow	45.90 (26.26)	59.77 (34.00)	52.08 (23.86)	51.04 (21.70)	50.82 (18.44)
P value†	p≤0.01**	0.45	0.27	0.75	0.07
Household size					
<5	58.64 (21.32)	58.77 (24.99)	51.16 (24.87)	49.19 (18.51)	55.32 (15.12)
5–10	61.82 (23.41)	62.16 (24.53)	58.16 (23.42)	54.76 (19.96)	59.84 (17.22)
11–15	66.31 (24.29)	70.51 (24.86)	66.67 (19.28)	59.90 (18.02)	65.83 (17.75)
>15	74.31 (21.18)	76.74 (15.28)	68.06 (20.26)	64.35 (9.82)	71.78 (13.61)
P value†	p≤0.05*	p≤0.01**	p≤0.01**	p≤0.01**	p≤0.01**
Location					
Urban	64.39 (22.78)	67.20 (21.87)	61.39 (23.77)	56.60 (18.94)	62.69 (16.78)
Rural	61.11 (23.19)	60.83 (25.84)	55.90 (23.44)	53.27 (19.32)	58.61 (17.02)
P value	0.25	p≤0.05	0.07	0.17	0.06
Literacy					
Illiterate	60.58 (22.96)	61.56 (25.70)	54.74 (25.63)	52.85 (20.52)	58.22 (17.37)
Literate	64.43 (23.06)	65.19 (23.12)	61.79 (20.55)	56.50 (17.42)	62.42 (16.37)
P value	0.172	0.228	p≤0.05*	0.121	p≤0.05*
Households' level of income (US\$)					

Continued

**Table 4** Continued

Socioeconomic characteristics	Emotional/information support	Tangible support	Positive interaction	Affectionate	Overall support index
<250	59.27 (22.63)	61.44 (25.01)	56.41 (22.79)	52.21 (19.39)	57.69 (16.30)
250–500	65.66 (23.68)	65.55 (22.87)	60.98 (23.75)	57.22 (18.41)	63.03 (17.49)
501–750	69.79 (22.05)	60.42 (24.43)	57.64 (28.01)	59.72 (18.82)	63.98 (19.11)
>750	62.31 (22.11)	71.88 (28.50)	57.29 (25.07)	54.17 (20.64)	62.09 (15.58)
P value†	0.080	0.280	0.580	0.140	0.080
<b>House ownership</b>					
No	59.42 (23.20)	59.56 (26.69)	53.97 (24.28)	51.71 (19.16)	56.96 (17.00)
Yes	66.08 (22.40)	67.90 (20.77)	63.07 (21.91)	58.12 (18.76)	64.21 (16.22)
P value	p≤0.05*	p≤0.01**	p≤0.01**	p≤0.01**	p≤0.01**
<b>Main households' occupation</b>					
Agriculture	58.65 (24.33)	56.49 (22.93)	56.49 (22.93)	52.49 (19.78)	57.47 (18.15)
Trade	65.25 (22.88)	66.06 (20.28)	66.06 (20.28)	57.11 (18.31)	63.57 (16.24)
Labour	63.52 (21.21)	54.35 (25.54)	54.35 (25.54)	53.17 (19.33)	59.98 (16.48)
Others	56.37 (26.06)	55.45 (24.94)	55.45 (24.94)	55.77 (21.44)	57.39 (17.22)
Formal job	68.75 (21.42)	63.38 (20.62)	63.38 (20.62)	58.84 (16.66)	64.71 (15.81)
P value†	0.126	0.849	0.054	0.43	0.156

Note: Figures in parentheses are SDs.

\*p&lt;0.05, \*\*p&lt;0.01.

†F-test while the rest are t-test.

**Table 5** Results of logistic regression

Socioeconomic characteristics	Unadjusted OR (95% CI)	Adjusted OR (95% CI)
<b>Age</b>		
15–20	1	1
21–30	2.87 (1.49 to 5.54)**	2.26 (1.04 to 4.89)*
31–40	4.13 (1.85 to 9.22)**	3.56 (1.28 to 9.89)*
41–50	4.41 (2.01 to 9.64)**	4.28 (1.60 to 11.26)**
51–60	5.55 (1.76 to 17.52)**	6.17 (1.55 to 24.59)**
<b>Sex</b>		
Female	1	1
Male	3.14 (1.90 to 5.19)**	2.73 (1.49 to 4.98)**
<b>Marital status</b>		
Unmarried	1	1
Married	0.94 (0.55 to 1.58)	1.05 (0.53 to 2.08)
Widow	0.20 (0.06 to 0.67)**	0.17 (0.04 to 0.80)*
<b>Household size</b>		
<5	1	1
5–10	1.58 (0.89 to 2.79)	1.96 (0.97 to 3.98)
11–15	2.91 (1.21 to 7.02)*	2.49 (0.89 to 6.94)
>15	6.61 (1.76 to 24.87)**	5.69 (1.32 to 24.65)*
<b>Location</b>		
Urban	1	1
Rural	0.62 (0.37 to 1.02)	0.71 (0.39 to 1.31)
<b>Literacy</b>		
Illiterate	1	1
Literate	1.22 (0.75 to 1.98)	0.86 (0.47 to 1.59)
<b>Monthly income (US\$)</b>		
<250	1	1
250–500	1.78 (0.62 to 5.16)	1.03 (0.31 to 3.42)
501–750	1.67 (0.97 to 2.90)	3.13 (1.00 to 9.83)*
>750	2.60 (1.02 to 6.64)*	2.00 (1.11 to 3.60)*
<b>House ownership</b>		
No	1	1
Yes	2.256 (1.37 to 3.71)**	1.997 (1.10 to 3.60)*
Hosmer and Lemeshow test	$\chi^2=7.47$ , df=8, p=0.487	

\*p&lt;0.05, \*\*p&lt;0.01.

counterparts. This phenomenon also substantiated that the marriage is an important source of social support. Married patients had higher social support.<sup>39,41</sup> Additionally, the factor of household size among the patients was a significant determinant of social support. Our finding implied that in a large-sized household, the patient would receive more social support from the family. The previously mentioned characteristics are both demonstrated in the research conducted in China.<sup>38</sup>

TB is more prevalent among the individuals with a lower socioeconomic status.<sup>6</sup> This implies that economic status is also one of the factors that plays a role in perceived social support of patients with TB. The findings of the current study showed that patients from groups with higher income levels received more social support. Mardan is a central district and people from the northern part of the province usually migrate and live in kacha (local term used for houses that are made from mud) houses or in slums that result in very poor living conditions. Patients from these households usually have low social support. Furthermore, in line with the view that the factor of homelessness is significant, we have included a proxy determinant that is the factor of home ownership. Therefore, the findings of the study showed that the factor of home ownership may be positively associated with care and social support from family and community.

### Strengths and limitations of the study

MOS-SSS 19-item scale is an instrument to measure social support and provides its usefulness for future studies in this region of Pakistan. This is a first attempt to measure the social support extended to patients with TB in Pakistan. The findings of the study capture the perceived social support among the population of patients with TB; however, these are not the representatives of the national-level population. Moreover, the selected participants of the study were under treatment for the last 4 weeks, and there might be a difference in the social support at the start and at the end of the treatment. Therefore, future research will explore that how the social support level may impact the adherence of patients to TB treatment.

### CONCLUSION

The previous findings show that it is paramount to address the treatment of TB within the context of the care and social support provided to patients, in addition to standard medical procedure. This study indicated that patients with pulmonary TB experienced an overall low level of care and social support from their families, friends and communities. Considering this issue, patients of TB should be considered as an integral part of the population who need to receive both clinical and social care. In addition, the findings of this study imply the need for a coordinated and integrated approach including not only clinical services but also encouragement of strong family and community support throughout the treatment. Individuals such as the poor, widows, women and marginalised patients should be extended more social support. Additionally, effective counselling and knowledge dissemination about TB ought to be present in families affected by TB at micro level, as well as society at the macro level.

**Acknowledgements** We extend our sincere thanks to the two anonymous reviewers and editor of this journal for their invaluable comments and suggestions on the earlier drafts of the manuscript. We also thank the study participants for their participation in the study and Mr Howard Goldman for proofreading the manuscript.



**Contributors** SES designed and performed the data collection and the analysis of the results. MMA supervised all the work from the start to the end. SP wrote the manuscript with inputs from other two authors.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient consent for publication** Obtained.

**Ethics approval** The study was approved by the research ethical review committee of the Asian Institute of Technology, Thailand, under reference number RERC-2017/001. In addition, the data were collected only after receiving written consent from the respondents. Authors followed all standard protocols needed to conduct this study.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data are available upon reasonable request.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

#### ORCID iD

Shahab E Saqib <http://orcid.org/0000-0003-0326-5197>

#### REFERENCES

- World Health Organization. *World tuberculosis report*. Geneva, Switzerland: WHO, 2016. <http://apps.who.int/iris/bitstream/10665/250441/1/9789241565394-eng.pdf?ua=1>
- World Health Organization EMRO. Global plan to stop TB, 2011-2015, 2015. Available: <http://www.emro.who.int/pak/programmes/stop-tuberculosis.html> [Accessed 2 Mar 2017].
- World Bank-WDI. Incidence of tuberculosis, 2014. Available: [http://search.worldbank.org/quickview?view\\_url=http%3A%2F%2Fdatabanksearch.worldbank.org%2FDataSearch%2FLoadReport.aspx%3Fdb%3D2%26cntrycode%3D%26sercode%3DSH.TBS.INCD%26yrcode%3D](http://search.worldbank.org/quickview?view_url=http%3A%2F%2Fdatabanksearch.worldbank.org%2FDataSearch%2FLoadReport.aspx%3Fdb%3D2%26cntrycode%3D%26sercode%3DSH.TBS.INCD%26yrcode%3D) [Accessed 20 Mar 2016].
- World Health Organization. Pakistan: stop tuberculosis who, East Mediterranean region, Cairo, 2015. Available: <http://www.emro.who.int/pak/programmes/stop-tuberculosis.html>
- Boccia D, Hargreaves J, De Stavola BL, et al. The association between household socioeconomic position and prevalent tuberculosis in Zambia: a case-control study. *PLoS One* 2011;6:e20824.
- Potter JL, Inamdar L, Okereke E, et al. Support of vulnerable patients throughout TB treatment in the UK. *J Public Health* 2016;38:391-5.
- Khanal S, Eisey H, King R, et al. Development of a patient-centred, psychosocial support intervention for multi-drug-resistant tuberculosis (MDR-TB) care in Nepal. *PLoS One* 2017;12:e0167559.
- Munro SA, Lewin SA, Smith HJ, et al. Patient adherence to tuberculosis treatment: a systematic review of qualitative research. *PLoS Med* 2007;4:e238.
- Cazabon D, Alsdurf H, Satyanarayana S, et al. Quality of tuberculosis care in high burden countries: the urgent need to address gaps in the care cascade. *Int J Infect Dis* 2017;56:111-6.
- Centers for Disease Control and Prevention. Basic TB facts, 2016. Available: <http://www.cdc.gov/tb/topic/basics/default.htm> [Accessed 9 Jul 2016].
- Long NH, Johansson E, Diwan VK, et al. Fear and social isolation as consequences of tuberculosis in Vietnam: a gender analysis. *Health Policy* 2001;58:69-81.
- Jittimanee SX, Nateniyom S, Kittikraisak W, et al. Social stigma and knowledge of tuberculosis and HIV among patients with both diseases in Thailand. *PLoS One* 2009;4:e6360.
- von dem Knesebeck O. Concepts of social epidemiology in health services research. *BMC Health Serv Res* 2015;15:357.
- Holt-Lunstad J, Smith TB, Layton JB. Social relationships and mortality risk: a meta-analytic review. *PLoS Med* 2010;7:e1000316.
- Liefoghe R, Michiels N, Habib S, et al. Perception and social consequences of tuberculosis: a focus group study of tuberculosis patients in Sialkot, Pakistan. *Soc Sci Med* 1995;41:1685-92.
- Mohammed S, Sajun SZ, Khan FS. Harnessing Photovoice for tuberculosis advocacy in Karachi, Pakistan. *Health Promot Int* 2015;30:262-9.
- House JS, Robbins C, Metzner HL. The association of social relationships and activities with mortality: prospective evidence from the Tecumseh community health study. *Am J Epidemiol* 1982;116:123-40.
- Harvey IS, Alexander K. Perceived social support and preventive health behavioral outcomes among older women. *J Cross Cult Gerontol* 2012;27:275-90.
- Sherbourne CD, Stewart AL. The mos social support survey. *Soc Sci Med* 1991;32:705-14.
- Paz-Soldán VA, Alban RE, Jones CD, et al. The provision of and need for social support among adult and pediatric patients with tuberculosis in Lima, Peru: a qualitative study. *BMC Health Serv Res* 2013;13:290.
- Nachega JB, Knowlton AR, Deluca A, et al. Treatment supporter to improve adherence to antiretroviral therapy in HIV-infected South African adults. A qualitative study. *J Acquir Immune Defic Syndr* 2006;43:S127-33.
- Government of Pakistan. *District at a glance*. Islamabad: Government of Pakistan, 2010.
- Government of Khyber Pakhtunkhwa. Mardan district demographics (census, 1998), 2017. Available: <http://kp.gov.pk/page/mardandistrictdemographics> [Accessed 2 Mar 2017].
- DHIS. *Disease pattern in out patient department. district health information system*, K.P.K. Peshawar: Department of Health, 2014.
- Government of Khyber Pakhtunkhwa. *The KP tuberculosis notification bill*. Peshawar: Government of K.P.K, 2016. <http://www.pakp.gov.pk/2013/bills/the-khyber-pakhtunkhwa-tuberculosis-notification-bill-2016/>
- District TB Office. *Consolidated report: consolidated report of TB reporting centers*. Mardan, 2016.
- Naing L, Winn T, Rusli B. Practical issues in calculating the sample size for prevalence studies. *Archives of orofacial Sciences* 2006;1:9-14.
- RAND. Social support survey instrument. Available: [https://www.rand.org/health/surveys\\_tools/mos/social-support/survey-instrument.html](https://www.rand.org/health/surveys_tools/mos/social-support/survey-instrument.html)
- Santos LM, Amorim LDAF, Santos DN, et al. Measuring the level of social support using latent class analysis. *Soc Sci Res* 2015;50:139-46.
- Gómez-Campelo P, Pérez-Moreno EM, de Burgos-Lunar C, et al. Psychometric properties of the eight-item modified medical outcomes study social support survey based on Spanish outpatients. *Qual Life Res* 2014;23:2073-8.
- Wang W, Zheng X, He H-G, et al. Psychometric testing of the Chinese mandarin version of the medical outcomes study social support survey in patients with coronary heart disease in mainland China. *Qual Life Res* 2013;22:1965-71.
- Alonso Fachado A, Montes Martínez A, Menendez Villalva C, et al. [Cultural adaptation and validation of the Medical Outcomes Study Social Support Survey questionnaire (MOS-SSS)]. *Acta Med Port* 2007;20:525-34.
- World Health Organization. *Management of substance abuse: process of translation and adaptation of instruments*. Geneva Switzerland: WHO, 2019. [https://www.who.int/substance\\_abuse/research\\_tools/translation/en/#](https://www.who.int/substance_abuse/research_tools/translation/en/#)
- Karim RM, Abdullah MS, Rahman AM, et al. Identifying role of perceived quality and satisfaction on the utilization status of the community clinic services; Bangladesh context. *BMC Health Serv Res* 2016;16:204.
- Getahun B, Nkosi ZZ. Satisfaction of patients with directly observed treatment strategy in Addis Ababa, Ethiopia: a mixed-methods study. *PLoS One* 2017;12:e0171209.
- Saqib SE, Ahmad MM, Amezcua-Prieto C, et al. Treatment delay among pulmonary tuberculosis patients within the Pakistan national tuberculosis control program. *Am J Trop Med Hyg* 2018;99:143-9.
- Saqib SE, Ahmad MM, Amezcua-Prieto C. Economic burden of tuberculosis and its coping mechanism at the household level in Pakistan. *Soc Sci J* 2018;55:313-22.
- Chen B, Peng Y, Zhou L, et al. Social support received by multidrug-resistant tuberculosis patients and related factors: a cross-sectional study in Zhejiang Province, People's Republic of China. *Patient preference and adherence* 2016;10:1063.
- Conte KP, Schure MB, Goins RT. Correlates of social support in older American Indians: the native elder care study. *Aging Ment Health* 2015;19:835-43.
- Ali TS, Krantz G, Gul R, et al. Gender roles and their influence on life prospects for women in urban Karachi, Pakistan: a qualitative study. *Glob Health Action* 2011;4:7448.
- Amare M, Hohfeld L, Jitsuchon S, et al. *Rural-Urban migration and employment quality: a case study from Thailand*. Asian development bank economics working paper series No.309, 2012.