

Provision of primary care by specialist physicians: a systematic review

Muhammad Jawad Hashim 

To cite: Hashim MJ. Provision of primary care by specialist physicians: a systematic review. *Fam Med Com Health* 2020;**8**:e000247. doi:10.1136/fmch-2019-000247

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/fmch-2019-000247>).

ABSTRACT

Patients with stable chronic diseases such as diabetes and hypertension can be safely managed at the primary care level. Yet many such patients continue to follow-up with specialists at a higher expense with no added benefit. We introduce a new term to describe this phenomenon: scope inversion, defined as the provision of primary care by specialist physicians. We aimed to quantify the extent of scope inversion by conducting a systematic review. MEDLINE and five other databases were searched using the keywords 'specialist AND (routine OR primary) AND provi*' as well as other variations. The search was limited to human research without restrictions on language or date of publication. The inclusion criterion was studies on rates of the provision of routine primary care by specialist physicians. Thirteen observational studies met the inclusion criteria. A wide range of primary care involvement was observed among specialists, from 2.6% to 65% of clinic visits. Among children, 41.3% of visits with specialists were routine follow-ups for conditions such as allergic rhinitis and seborrhoeic dermatitis which could be managed in primary care. Data quality was moderate to low across the studies due to limitations of source data and varying definitions of primary care. Specialist physicians provide primary care to patients in a substantial proportion of clinic visits. Scope inversion is wasteful as it diverts patients to more expensive care without improving outcomes. A systems approach is needed to mitigate scope inversion and its harmful effects on healthcare service delivery.

INTRODUCTION

Common medical conditions, which can be safely managed in primary care, are sometimes treated by specialist physicians.¹ A new term is introduced in this article to highlight this health systems issue: scope inversion is defined as the provision of primary care by specialist physicians. It can occur as routine follow-up of chronic diseases or management of mild acute illnesses by a specialist (figure 1). Health systems aim to triage patients to appropriately qualified professionals. Scope inversion is inappropriate triage. As a health systems problem, scope inversion needs identification, quantification and mitigation.

Scope inversion can undermine the healthcare system in several insidious ways. Primary care provision by subspecialists leads to more expensive care without improving outcomes.

Tertiary care clinics can be bogged down by the large number of patients with primary care needs thus resulting in longer waiting times for available appointments for patients who genuinely need specialist care.² For example, patients who need appointments for specialty care can face waiting times of over 40 days.³ Patients may expect to receive all of their care from specialists including treatment of common illnesses, further undermining primary care and wasting tertiary care resources. Specialist physicians do not perform the full scope of primary care such as preventive services and care coordination for patients with multiple chronic diseases.⁴ Quality measures based on specialist-oriented guidelines are often inappropriately applied to primary care settings. The value of primary care extends to the care of the whole person and the overall health of populations.⁵

Although studies on referrals by family physicians and general practitioners (GPs) to specialists have been published repeatedly,⁶ there is limited research on the issue of patients returning to GPs to resume continuity of care. There is some evidence that in health systems with robust primary care, generalists provide more efficient care.⁷ For instance, medical expenses per episode of back pain are almost twice as high with orthopaedic surgeons compared with primary care physicians (PCPs).⁸ Specialists are likely to utilise more diagnostic tests and procedures.⁸ In health services research, care by specialist physicians does not result in better outcomes such as time to recovery or control of chronic diseases.⁹

Thus, the provision of primary care by specialists can undermine healthcare delivery in numerous ways. As this phenomenon is relatively understudied, we aimed to quantify the extent of scope inversion by conducting a systematic review.

METHODS

Definitions

A generalist may be defined as a physician (or other healthcare provider such as a



© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

Department of Family Medicine, United Arab Emirates University, Al Ain, United Arab Emirates

Correspondence to

Dr Muhammad Jawad Hashim; jhashim@uaeu.ac.ae, physicianthinker@gmail.com




Figure 1 Scope inversion: the provision of primary care by specialist physicians.

nurse practitioner) trained to manage a wide scope of illnesses, whereas ‘specialist’ denotes a clinician who is trained to manage a narrow range of medical conditions (eg, a nephrologist or an ophthalmologist). A PCP is a generalist who provides continuity of care to patients while coordinating care with specialists. Such physicians include general physicians/internists, general paediatricians and GPs. The terms GP and family physician are used synonymously in the literature to indicate a PCP who has received training in the clinical specialty of General Practice/Family Medicine.⁵ General Practice/Family Medicine is defined as the medical specialty that manages common and long-term illnesses in children and adults focusing on overall health and well-being.¹⁰

Search strategy

A systematic review was conducted using Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines.¹¹ The PRISMA statement provides a framework for conducting a comprehensive and reproducible review of research studies on a topic. MEDLINE, EBSCO Academic Search Complete, Scopus, ScienceDirect, Web of Science and Embase databases were searched using the keywords ‘specialist AND (routine OR primary) AND provi*’ as well as other variations (online supplementary appendix). The inclusion criterion was studies on rates of provision of routine primary care by specialist physicians. All study designs were considered. Exclusion criteria were review articles with no original data, and articles on shared comanagement of patients by specialists and generalists, or on referrals from GPs to specialists. A trained medical research librarian experienced in conducting systematic reviews provided guidance on search strategy and methodology. The search was conducted over several months with the final refresh in December 2019.

Search results were screened for appropriate articles based on the titles and abstracts initially (see the online supplementary appendix for details on study selection). It was not feasible to combine the results from different studies in a meta-analysis due to the heterogeneity of individual study designs. Data quality was moderate to low across the studies due to limitations of source data and varying definitions of primary care.




Figure 2 Estimates of scope inversion.

RESULTS

A wide range of estimates were found in the 13 studies that met inclusion criteria (Figure 2 and online supplementary appendix). An early study found the involvement of specialists in primary care to be between 9% and 34% depending on the definition of primary care.¹² Data from the US National Ambulatory Medical Care Survey (NAMCS) in 2002 showed that in about one-half of all clinic visits, specialist physicians indicated that they were the PCP for the patient.¹³ Among children, 41.3% of visits with specialists were routine follow-ups for conditions such as allergic rhinitis and seborrhoeic dermatitis which ‘could be managed in primary care settings’.¹⁴ In the 2013 NAMCS, general surgeons, for instance, reported that they were providing primary care in 2.6% of patient visits (table 1; NAMCS 2013 data).¹⁵ According to a US report on the care of the elderly, only 60% identified a family physician as their usual source of healthcare.¹⁶ The remaining 40% of respondents were potentially receiving primary care from specialists. In a US population-based survey, 14% of respondents named a specialist as their source of primary care.¹⁷ For example, 6% reported receiving primary healthcare from a cardiologist.¹⁷ In fact, about half of the respondents were potentially relying on a specialist for their routine medical needs.¹⁷ A review of earlier NAMCS data found that in 44% of visits to nephrologists and 65% of visits to infectious disease consultants, the specialist physicians reported being the PCP.⁹ In Colorado, 46.5% of 2745 specialists reported providing primary care.¹⁸ A retrospective analysis of a large US Medicare database reported that 25% of beneficiaries received care from a specialist.¹⁹

DISCUSSION

Specialist physicians provide primary care to patients in a substantial proportion of clinic visits, ranging from 2.6% to 65% of their consultations. This variation may be explained by the heterogeneity of study designs, data quality, different ways of defining primary care and the level of involvement of specialists. The latter has been

Table 1 Provision of primary care by specialists, according to specialty (n=2879)

Specialist's field	Percent of office visits identified by the specialist as a visit to the PCP* (SE)
Cardiology	10.0 (3.3)
Psychiatry	8.0 (5.3)
Otolaryngology	5.6 (4.5)
General surgery	2.6 (1.3)
Ophthalmology	2.0 (1.0)
Orthopaedic surgery	1.0 (0.5)
Other specialists	8.1 (1.4)

Source: National Ambulatory Medical Care Survey,¹⁵

*PCP is the patient's primary care provider as indicated by a positive response to the question, 'Are you the patient's primary care physician/provider?'

categorised in a typology by Kendall *et al*: family physician-dominated comanagement, specialist-dominated comanagement and exclusively specialist care.²⁰ In contrast, multidisciplinary shared care occurs when GPs refer patients to allied health professionals such as dieticians, physical therapists and psychologists.²¹

We know that specialists provide less comprehensive primary care than generalists such as family physicians.⁹ Thus, specialists tend to see fewer new problems, treat a limited range of conditions and provide less preventive care.⁷ The scope of practice among specialist physicians is narrower in terms of the range of illnesses seen.²² Although patients may identify a gynaecologist as their 'PCP', they may not be receiving comprehensive primary care.²³ Primary care in this context is more than the care of a chronic condition. A specialist might counsel a patient to quit smoking if it helps to improve the disease status; however, this should not be construed as full primary care.

It is well recognised that a case mix of patients with the mild disease is well suited to the continuity of care by primary care doctors. For specialist physicians, having these stable patients on their panel improves performance measures while avoiding high-risk cases.²⁴ By shifting to stable patients with the controlled disease, specialists avoid clinical care that is more time and effort-intensive. Among patients with stable chronic diseases, routine office visits are less complex and often billed at the same level as those with complex, unstable problems. Financial inducement may play a role in the 'perennial follow-up' of stable patients by specialists.²⁵

The care provided by generalists and specialists is similar to common chronic diseases.^{26,27} Patients perceive family physicians positively when transitioning back from tertiary to primary care.²⁸ Thus, there seems to be no evidence-based rationale to continue the follow-up of patients with the mild stable disease with subspecialists. It is not surprising that the likelihood of consultation with a specialist physician by elderly patients is more linked to

their incomes than healthcare needs.²⁹ Gaps in specialists' clinical knowledge base often lead to cross-referrals to other specialties resulting in tortuous patient journeys leading to diagnostic delays.³⁰

Some experts have noted that the shortage of family physicians may necessitate the provision of primary care by specialists.^{9,12,31} As such, continuity of care provided by these specialists is better than having no usual source of care.³² Examples include ongoing care of HIV+ patients by infectious disease specialists and the care of patients on dialysis by nephrologists.²⁰ However, these stopgap measures should not be viewed as a long-term substitute for accessible and affordable primary care.

Limitations of this review include a dearth of studies, especially the lack of published research from outside the USA. The varying definitions of primary care and metrics used to quantify it make it difficult to pool the results in a meta-analysis or produce a summary statistic. Specialist physicians may gradually switch into generalist disciplines later in their careers.³³ The extent of primary care provision by specialists seems to have changed over the last few decades.⁹

Potential solutions to reduce scope inversion

The return of patients from specialists to primary care can be a potential 'fault line' in the continuum of care.³⁴ This observation underscores the need for a clear summary (such as a hospital discharge note) and a detailed patient care plan to be provided by the specialist to the GP.³⁵ Interclinician communication can guide the transfer of patient management from specialty to primary care.³⁶ Family physicians have raised this concern in earlier research.³⁷ Discussion of the care plan between the specialist and the family physician, either face-to-face or via teleconferencing, can reduce hospital readmissions and improve independent living and quality of life.³⁸

Reminder letters to patients have improved follow-up rates with GPs from a baseline of 56%–90% after the mailings.³⁹ Innovative solutions such as a telemedicine service can reduce scope inversion by decreasing the need for face-to-face consultations with a specialist.⁴⁰ Community health services can augment the care provided by family physicians without transferring to a specialist.⁴¹ Such strategies underscore the importance of the safe and timely return of patients to the care of community-based GPs.³⁷ Generalists are essential to guide patients given the rising multimorbidity of chronic illnesses.⁴² The overall goal of healthcare is integrated primary-specialist care exemplified by teamwork, information exchange, shared guidelines, continuous training, access to care and sustainable financing. Enforcement of timely return of stable patients to primary care is perhaps needed to reduce scope inversion.

Acknowledgements The authors would like to thank Linda Ostlundh, research librarian and director of the National Medical Library for her guidance in conducting the searches for the systematic review.

Contributors MJH: conceived of the research idea, conducted the literature search and wrote the manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Muhammad Jawad Hashim <http://orcid.org/0000-0002-9280-9709>

REFERENCES

- Starfield B, Lemke KW, Herbert R. Comorbidity and the use of primary care and specialist care in the elderly. *Ann Fam Med* 2005;3:215–22.
- Largey G, Chakraborty S, Tobias T, et al. Audit of referral pathways in the diagnosis of lung cancer: a pilot study. *Aust J Prim Health* 2015;21:106–10.
- Baume PE. Waiting times for non-urgent specialist appointments. *Med J Aust* 1995;162:648–9.
- Woolhandler S, Himmelstein DU. Administrative work consumes one-sixth of U.S. physicians' working hours and lowers their career satisfaction. *Int J Health Serv* 2014;44:635–42.
- Hashim MJ. Principles of family medicine and general practice – defining the five core values of the specialty. *J Prim Health Care* 2016;8:283–7.
- Lin F, Sun Q, Peng Z, et al. The innovations in China's primary health care reform: Development and characteristics of the community health services in Hangzhou. *Fam Med Community Health* 2015;3:52–66.
- Rosenblatt RA, Hart LG, Baldwin LM, et al. The generalist role of specialty physicians: is there a hidden system of primary care? *JAMA* 1998;279:1364–70.
- Harrold LR, Field TS, Gurwitz JH. Knowledge, patterns of care, and outcomes of care for generalists and specialists. *J Gen Intern Med* 1999;14:499–511.
- Edwards ST, Mafi JN, Landon BE. Trends and quality of care in outpatient visits to generalist and specialist physicians delivering primary care in the United States, 1997–2010. *J Gen Intern Med* 2014;29:947–55.
- Jawad Hashim M, Hashim MJ. A definition of family medicine and general practice. *J Coll Physicians Surg Pak* 2018;28:76–7.
- Moher D, Liberati A. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann Intern Med* 2009;151:264–9.
- Spiegel JS, Rubenstein LV, Scott B, et al. Who is the primary physician? *N Engl J Med* 1983;308:1208–12.
- Woodwell DA, Cherry DK. National ambulatory medical care survey: 2002 summary. *Adv Data* 2004;346:1–44.
- Valderas JM, Starfield B, Forrest CB, et al. Routine care provided by specialists to children and adolescents in the United States (2002–2006). *BMC Health Serv Res* 2009;9:221.
- National Center for Health Statistics. National Ambulatory Medical Care Survey: 2013 State and National Summary Tables, 2013. Available: https://www.cdc.gov/nchs/data/ahcd/namcs_summary/2013_namcs_web_tables.pdf [Accessed 13 Dec 2016].
- Mold JW, Fryer GE, Phillips RL, et al. Family physicians are the main source of primary health care for the Medicare population. *Am Fam Physician* 2002;66:2032.
- Stock Keister MC, Green LA, Kahn NB, et al. Few people in the United States can identify primary care physicians. *Am Fam Physician* 2004;69:2312.
- Fryer GE, Consoli R, Miyoshi TJ, et al. Specialist physicians providing primary care services in Colorado. *J Am Board Fam Pract* 2004;17:81–90.
- Romaire MA, Haber SG, Wensky SG, et al. Primary care and specialty providers: an assessment of continuity of care, utilization, and expenditures. *Med Care* 2014;52:1042–9.
- Kendall CE, Younger J, Manuel DG, et al. Health administrative data can be used to define a shared care typology for people with HIV. *J Clin Epidemiol* 2015;68:1301–11.
- Chan B, Proudfoot J, Zwar N, et al. Satisfaction with referral relationships between general practice and allied health professionals in Australian primary health care. *Aust J Prim Health* 2011;17:250–8.
- Freeman J, Petterson S, Bazemore A. Accounting for complexity: aligning current payment models with the breadth of care by different specialties. *Am Fam Physician* 2014;90:790.
- Fink KS, Baldwin LM, Lawson HW, et al. The role of gynecologists in providing primary care to elderly women. *J Fam Pract* 2001;50:153–8.
- Casalino LP, Alexander GC, Jin L, et al. General internists' views on pay-for-performance and public reporting of quality scores: a national survey. *Health Aff* 2007;26:492–9.
- Ackerman SL, Gleason N, Monacelli J, et al. When to Repatriate? Clinicians' Perspectives on the Transfer of Patient Management from Specialty to Primary Care. *J Gen Intern Med* 2014;29:1355–61.
- Smetana GW, Landon BE, Bindman AB, et al. A comparison of outcomes resulting from generalist vs specialist care for a single discrete medical condition: a systematic review and methodologic critique. *Arch Intern Med* 2007;167:10–20.
- Mendenhall RC, Moynihan CJ, Radecki SE. The relative complexity of primary care provided by medical specialists. *Med Care* 1984;22:987–1001.
- Dutton H, Rowan MS, Liddy C, et al. Patient perspectives on discharge from specialist type 2 diabetes care back to primary care: a qualitative study. *Can J Diabetes* 2014;38:191–7.
- Kaplan MS, Huguet N, Feeny D, et al. Visits to family physicians or specialists by elderly persons in Canada and the United States: an exploratory comparative study. *Int J Health Serv* 2010;40:139–49.
- Scott IA, Campbell DA. The gaps in specialists' diagnoses. *Med J Aust* 2018;208:196–7.
- Aiken LH, Lewis CE, Craig J, et al. The contribution of specialists to the delivery of primary care. *N Engl J Med* 1979;300:1363–70.
- Spatz ES, Sheth SD, Gosch KL, et al. Usual source of care and outcomes following acute myocardial infarction. *J Gen Intern Med* 2014;29:862–9.
- Christakis NA, Jacobs JA, Messikomer CM. Change in self-definition from specialist to generalist in a national sample of physicians. *Ann Intern Med* 1994;121:669–75.
- Mitchell GK. Improving coordination of care between specialist and general practice for people with chronic pain. *Med J Aust* 2013;198:301–2.
- Corwin P, Bolter T. The effects of audit and feedback and electronic referrals on the quality of primary care referral letters. *J Prim Health Care* 2014;6:324–7.
- Ackerman SL, Gleason N, Monacelli J, et al. When to repatriate? Clinicians' perspectives on the transfer of patient management from specialty to primary care. *J Gen Intern Med* 2014;29:1355–61.
- Brez S, Rowan M, Malcolm J, et al. Transition from specialist to primary diabetes care: a qualitative study of perspectives of primary care physicians. *BMC Fam Pract* 2009;10:39.
- Mitchell GK. End-of-life care for patients with cancer. *Aust Fam Physician* 2014;43:514–9.
- Foley F, Selzer R, McGartland M. Improving patient return to general practice after specialist assessment. *Australas Psychiatry* 2011;19:236–9.
- Muir J. Telehealth: the specialist perspective. *Aust Fam Physician* 2014;43:828–30.
- Young CE, Mutch AJ, Boyle FM, et al. Investigating referral pathways from primary care to consumer health organisations. *Aust J Prim Health* 2010;16:260–7.
- Warner DF, Schiltz NK, Stange KC, et al. Complex multimorbidity and health outcomes in older adult cancer survivors. *Fam Med Community Health* 2017;5:129–38.